

Limit superior and limit inferior:

$$a_n = (-1)^n + \frac{1}{n} \quad n \geq 1$$


$$a_1 = 0 \quad a_2 = 3/2 = 1.5$$

$$a_3 = -2/3 = -0.66 \quad a_4 = 5/4 = 1.25$$

$$a_5 = -4/5 = -0.8 \quad a_6 = 7/6 = 1.16$$

$$a_7 = -6/7 = -0.85 \quad a_8 = 9/8 = 1.12$$

$$a_9 = -8/9 = -0.88 \quad a_{10} = 11/10 = 1.1$$



stuck at -1

moves till 1.

Claim: Let $(a_n)_{n \geq m}$ be a sequence of real

numbers. Let

$$L^+ = \limsup_{n \rightarrow \infty} (a_n) \in \mathbb{R} \cup \{\pm \infty\}$$

$$L^- = \liminf_{n \rightarrow \infty} (a_n) \in \mathbb{R} \cup \{\pm \infty\}.$$

① For every $x > L^+$, $\exists N \geq m$ s.t.

$$x > a_n \quad \forall n \geq N.$$

② For every $y < L^-$, $\exists N \geq m$ s.t.

$$y < a_n \quad \forall n \geq N.$$

Proof: Given $x > L^+$

$$\Rightarrow x > \inf_{n \geq m} (a_n^+)$$

Since $x > \inf_{n \geq m} (a_n^+)$, there must be

at least one N_0 s.t. $x > a_{N_0}^+$

$$\text{if not then } x \leq a_{N_0}^+ \quad \forall N_0 \geq m$$

$\Rightarrow x$ is a lower bound on $(a_n^+)_{n \geq m}$ that

is greater than $\inf_{n \geq m} (a_n^+) \Rightarrow x$ is infimum.

$$\text{Now } x > a_{N_0}^+ = \sup_{n \geq N_0} (a_n)$$

$$\Rightarrow x > a_n \quad \forall n \geq N_0.$$

② $y < L^- = \sup_{n \geq m} (a_n^-)$

$$\Rightarrow \exists N_0 \geq m \text{ s.t. } y < a_{N_0}^-$$

$$\text{otherwise } y \geq a_n^- \quad \forall n \geq m$$

$$\Rightarrow y \text{ is a supremum}$$

That is $\exists N_0$ $y < a_{N_0}^-$

$$= \inf_{n \geq N_0} (a_n)$$

$$\Rightarrow y < a_n \quad \forall n \geq N_0.$$

$$\textcircled{3} \quad \inf_{n=m}^{\infty} (a_n) \leq L^- \leq L^+ \leq \sup_{n=m}^{\infty} (a_n)$$

Proof:

$$\inf_{n=m}^{\infty} (a_n) = a_m^- \leq a_{m+1}^- \leq \dots \quad \text{For all } N \geq m$$

$$\Rightarrow \inf_{n \geq m} (a_n) \leq a_N^-$$

$$= \inf_{n \geq N} (a_n)$$

$$\leq a_N \quad \forall N \geq m$$

$$\leq \sup_{n \geq N} (a_n) = a_N^+$$

$$\leq \sup_{n=m}^{\infty} (a_n)$$

$$\Rightarrow \inf_{n \geq m} (a_n) \leq a_N^- \leq a_N^+ \leq \sup_{n=m}^{\infty} (a_n)$$

Taking $\lim_{N \rightarrow \infty} \Rightarrow$

$$\inf_{n \geq m} (a_n) \leq L^- \leq L^+ \leq \sup_{n=m}^{\infty} (a_n).$$

④ If c is any limit point of $(a_n)_{n \geq m}$

$$\text{then } L^- \leq c \leq L^+.$$

Proof: $\forall N \geq m \quad \exists n \geq N$ s.t.

$$|a_n - c| \leq \varepsilon \quad \forall \varepsilon > 0.$$

$$\text{or } c - \varepsilon \leq a_n \leq c + \varepsilon \quad \forall \varepsilon > 0.$$

$$\Rightarrow c \leq a_n \leq c$$

$$\text{Now } a_n^+ = \sup_{n \geq N} (a_n)$$

$$\geq a_{N_0} \geq c$$

$$\lim_{N \rightarrow \infty} a_n^+ \geq \lim_{N \rightarrow \infty} c$$

$$\Rightarrow L^+ \geq c.$$

$$a_n^- = \inf_{n \geq N} (a_n)$$

$$\leq a_{N_0} \leq c$$

$$\Rightarrow L^- \leq c$$

$$\text{Combining } L^- \leq c \leq L^+.$$

⑤ If L^+ is finite then it is a limit point of $(a_n)_{n=m}^{\infty}$. Similarly, L^- is limit point of finite.

Proof:

Let $\varepsilon > 0$ and $N \geq m$ be arbitrary.

$$\text{define } a_N^+ = \sup_{n=N}^{\infty} (a_n) \quad (N \geq m)$$

$$\text{Since } \lim_{N \rightarrow \infty} a_N^+ = L^+$$

$$\exists N_0 \text{ s.t. } \forall K \geq N_0$$

$$|a_K^+ - L^+| \leq \varepsilon/2 \quad \forall \varepsilon > 0$$

$$\Rightarrow a_K^+ \leq L^+ + \varepsilon/2$$

Define

$$K = \max(M, N_0)$$

$$\Rightarrow K \geq N_0 \Rightarrow a_K^+ \leq L^+ + \varepsilon/2$$

$$\Rightarrow a_K^+ - \varepsilon/2 \leq L^+$$

$$\text{Since } a_K^+ = \sup_{n \geq K} (a_n)$$

$$\Rightarrow a_K^+ - \varepsilon/2 \text{ can't be supremum.}$$

$$\Rightarrow \exists n \geq K \text{ s.t. } a_n > a_K^+ - \varepsilon/2$$

$$L^+ - \varepsilon/2 \leq a_K^+ < a_n + \varepsilon/2 \leq a_K^+ + \varepsilon/2$$

$$\leq L^+ + \varepsilon$$

$$< L^+ + 3\varepsilon/2$$

$$\Rightarrow L^+ - \varepsilon < a_n < L^+ + \varepsilon$$

$\Rightarrow$ We have found $n \geq K \geq M$ s.t.

$$|a_n - L^+| < \varepsilon. \text{ This is true for all } \forall \varepsilon > 0.$$

Same for the other case.

⑥ Let c be a real number. If $(a_n)_{n \geq m}$

converges to c then $L^+ = c = L^-$. Converse is

true as well.

Proof:

Let $\lim_{n \rightarrow \infty} a_n = c$ or

$$\forall \varepsilon > 0 \quad \exists N \text{ s.t. } \forall n \geq N$$

$$|a_n - c| \leq \varepsilon.$$

$$\Rightarrow c - \varepsilon \leq a_n \leq c + \varepsilon.$$

$$\sup_{n \geq N} (a_n) \leq c + \varepsilon \quad [\text{supremum can be smaller than upper bound}]$$

$$\text{or } a_N^+ \leq c + \varepsilon$$

$$\text{similarly } \inf_{n \geq N} (a_n) \geq c - \varepsilon \quad [\text{infimum can be larger than lower bound}]$$

$$\Rightarrow L^+ \leq c, \quad L^- \geq c$$

$$\Rightarrow L^- \geq c \geq L^+ \quad \text{but } L^- \leq L^+$$

$$\Rightarrow L^+ = L^- \Rightarrow L^+ = L^- = c.$$

$$\Rightarrow \text{let } L^+ = L^- = c$$

$$\Rightarrow \lim_{n \rightarrow \infty} a_n^+ = c$$

$$\exists N_0 \text{ s.t. } \forall n \geq N_0$$

$$|a_n^+ - c| \leq \varepsilon \quad \forall \varepsilon > 0.$$

$$\exists N_1 \text{ s.t. } \forall n \geq N_1$$

$$|a_n^+ - c| \leq \varepsilon \quad \forall \varepsilon > 0$$

$$\text{Take } N = \max\{N_0, N_1\}$$

$$c - \varepsilon \leq a_N^- \leq a_n \leq a_N^+ \leq c + \varepsilon \quad \forall n \geq N$$

$$\Rightarrow |a_n - c| \leq \varepsilon \quad \forall n \geq N$$

$$\Rightarrow (a_n) \text{ converges to } c.$$

S Lecture 12.1

Thursday, 18 September 2025 22:20

Claim: (a_n) (b_n) be two sequences of

real numbers. Then $a_n \leq b_n \quad \forall n \geq m$

imply:

$$\sup(a_n) \leq \sup(b_n)$$

$$\inf(a_n) \leq \inf(b_n)$$

$$\limsup(a_n) \leq \limsup(b_n)$$

$$\liminf(a_n) \leq \liminf(b_n).$$

Proof: ① $\forall n \geq m \quad a_n \leq b_n \leq \sup(b_n)$

$$\Rightarrow \sup(b_n) \text{ is an upper bound on } (a_n)$$

$$\Rightarrow \sup(a_n) \leq \sup(b_n).$$

② $\inf(a_n) \leq a_n \leq b_n$

$$\Rightarrow \inf(a_n) \text{ is a lower bound on } (b_n)$$

$$\Rightarrow \inf(a_n) \leq \inf(b_n) \quad (\text{But greatest lower bound is } \inf(b_n))$$

③ $a_n^+ = \sup(a_n)_{n \geq N} \leq \sup(b_n)_{n \geq N} = b_n^+$

$$a_n^+ \leq b_n^+$$

$$\Rightarrow \inf a_n^+ \leq \inf b_n^+ \Leftrightarrow \limsup(a_n) \leq \limsup(b_n)$$

④ similarly.

Claim: Squeeze test.

Let (a_n) (b_n) (c_n) be sequences of real numbers. s.t.

$$a_n \leq b_n \leq c_n \quad \forall n \geq m.$$

If $\lim_{n \rightarrow \infty} a_n = L = \lim_{n \rightarrow \infty} b_n$ then $\lim_{n \rightarrow \infty} c_n = L$.

Proof: A priori it is not clear if (c_n) is

convergent. But if we can show that

$$L^+ \text{ of } (b_n) = L^- \text{ of } (b_n) \text{ then}$$

$$\lim_{n \rightarrow \infty} b_n = L.$$

$$a_n \leq b_n \leq c_n$$

$$\Rightarrow \liminf a_n \leq \liminf b_n \leq \liminf c_n$$

$$L \leq \liminf(b_n) \leq L$$

$$\Rightarrow \liminf(b_n) = L.$$

$$a_n \leq b_n \leq c_n$$

$$\Rightarrow \limsup(a_n) \leq \limsup(b_n) \leq \limsup(c_n)$$

$$L \leq \limsup(b_n) \leq L$$

$$\limsup(b_n) = L$$

$$\Rightarrow \lim_{n \rightarrow \infty} b_n = L.$$

Ex: Let $a_n = \frac{(-1)^n}{n} + \frac{1}{n^2} \quad n \geq 1$

$$-1+1=0$$

$$\frac{1}{2} + \frac{1}{4} = \frac{3}{4}$$

$$-\frac{1}{3} + \frac{1}{9} = -\frac{2}{9}$$

$$\frac{1}{4} + \frac{1}{16} = \frac{5}{16}$$

$$\frac{(-1)^n}{n} + \frac{1}{n^2} \leq \frac{1}{n} + \frac{1}{n} \cdot \frac{1}{n}$$

$$= \frac{1}{n} \left(\frac{1}{n} + 1 \right)$$

$$\leq \frac{2}{n}$$

$$\frac{(-1)^n}{n} + \frac{1}{n^2} \geq -\frac{1}{n} - \frac{1}{n} = -\frac{2}{n}$$

$$\Rightarrow -\frac{2}{n} \leq a_n \leq \frac{2}{n}$$

Then two converge to 0

$$\Rightarrow \lim_{n \rightarrow \infty} a_n = 0.$$

Or. $\bar{a}^n \leq \frac{1}{n}$

$$\text{For } n=1 \quad \frac{1}{2} \leq 1$$

$$\bar{a}^{(n+1)} = \frac{1}{2} \bar{a}^n \leq \frac{1}{2n}$$

$$\leq \frac{1}{n+1}$$

$$\Rightarrow 0 \leq \bar{a}^n \leq \frac{1}{n} \Rightarrow \lim_{n \rightarrow \infty} \bar{a}^n = 0.$$

Theorem: (Completeness of reals)

A sequence of real numbers is a Cauchy sequence

if and only if it is convergent.

Proof: Every convergent sequence is Cauchy, i.e.

proved already.

Now every Cauchy sequence is convergent in reals.

Let (a_n) be a Cauchy sequence. Thus

(a_n) is bounded.

$$\Rightarrow L^+ \text{ and } L^- \text{ are bounded.}$$

If $L^+ = L^-$ then the sequence converges.

Let $\varepsilon > 0$ be any real number. Since (a_n)

is Cauchy sequence $\Rightarrow$

$$\exists N \text{ s.t. } \forall m, n \geq N$$

$$|a_n - a_m| \leq \varepsilon \quad \forall \varepsilon > 0$$

$$\text{or } a_n - \varepsilon \leq a_n \leq a_n + \varepsilon \quad \forall n \geq N$$

$$\Rightarrow a_n - \varepsilon \leq \inf(a_n)_{n \geq k} \quad \forall k \geq N$$

$$\Rightarrow \lim_{k \rightarrow \infty} (a_n - \varepsilon) \leq \liminf(a_n)$$

$$\Rightarrow L^- \geq a_n - \varepsilon \rightarrow \textcircled{1}$$

$$\sup(a_n)_{n \geq k} \leq a_n + \varepsilon$$

$$\Rightarrow \lim_{k \rightarrow \infty} \sup(a_n)_{n \geq k} \leq a_n + \varepsilon$$

$$L^+ \leq a_n + \varepsilon \rightarrow \textcircled{2}$$

$$L^+ - L^- \leq a_n + \varepsilon - a_n + \varepsilon$$

$$= 2\varepsilon.$$

$$L^+ - L^- \geq 0 \geq -2\varepsilon$$

$$\Rightarrow |L^+ - L^-| \leq 2\varepsilon \quad \forall \varepsilon > 0$$

$$\Rightarrow L^+ - L^- = 0 \Rightarrow L^+ = L^-$$

$$\Rightarrow \text{sequence } (a_n) \text{ converges.}$$